

**WASHINGTON OB/GYN ASSOCIATES**

95 Leonard Avenue  
Building #1, Suite 300  
Washington, PA 15301  
724-225-3640 Phone

**McMURRAY OB/GYN ASSOCIATES**

2001 Waterdam Plaza  
Suite 201  
McMurray, PA 15317  
724-942-5420 Phone

**WASHINGTON OB/GYN ASSOCIATES**

**WAYNESBURG**  
343 East Roy Furman Highway  
Suite 104  
Waynesburg, PA 15370  
724-627-7928 Phone

**WASHINGTON OB/GYN ASSOCIATES**


**MON VALLEY-CHARLEROI**  
1200 McKean Avenue  
Charleroi, PA 15022  
724-483-1340 Phone

[www.WashingtonObGyn.net](http://www.WashingtonObGyn.net)

## What is group B streptococcus (beta strep)?

Group B streptococcus (also called beta strep, or GBS), is a very common type of bacteria. Ten to thirty percent of pregnant women carry GBS bacteria. Most often the bacteria are in the vagina or rectum. GBS bacteria are different from the type of bacteria that cause strep throat.

Healthy adults carrying GBS may not have any symptoms or problems. However, sometimes the bacteria can cause an infection in the uterus, bladder, kidneys, or, rarely, the brain (meningitis). These infections in adults are usually not serious and can be treated with antibiotics. But a baby can get very sick and even die if the mother has untreated GBS.



## How does it occur?

Both women and men may have GBS. The bacteria often live in the mouth, bowel, bladder, vagina, and rectum. A woman who carries GBS can pass it to her baby during labor or delivery.

## What are the symptoms?

In most cases pregnant women do not have any symptoms of infection. When they do have symptoms, they may include:

- fever over 100°F during labor
- painful, bloody, frequent, or urgent urination.

## How is it diagnosed?

GBS can be found by testing urine or swabbing the cervix, vagina, or rectum. The samples of fluid are cultured in a lab. A test result is positive if beta strep bacteria are found in the culture. The test is negative if the bacteria are not found. The Centers for Disease Control (CDC) recommends that all pregnant women have a GBS culture in the 35th to 37th weeks of pregnancy. A culture may also be done if you have premature rupture of the membranes, a fever during labor, or urinary tract symptoms such as painful, bloody, or frequent urination.

A rapid test for GBS may be done when you go to the hospital to deliver your baby. The results of this test can be known in 2 to 3 hours. However, it is less accurate than cultures and is rarely used earlier in a pregnancy.

## How is it treated?

If your tests for beta strep are positive, you will probably not be treated until you are in labor. During labor you will be given intravenous (IV) antibiotics. If you were treated earlier in the pregnancy, the bacteria could come back again before the baby is born. The one exception is when a urine culture, done at any time during pregnancy, shows that GBS is causing a bladder infection. In this case your healthcare provider will probably prescribe an oral antibiotic to treat the infection.

**WASHINGTON OB/GYN ASSOCIATES**

95 Leonard Avenue  
Building #1, Suite 300  
Washington, PA 15301  
724-225-3640 Phone

**McMURRAY OB/GYN ASSOCIATES**

2001 Waterdam Plaza  
Suite 201  
McMurray, PA 15317  
724-942-5420 Phone

**WASHINGTON OB/GYN ASSOCIATES**

**WAYNESBURG**  
343 East Roy Furman Highway  
Suite 104  
Waynesburg, PA 15370  
724-627-7928 Phone

**WASHINGTON OB/GYN ASSOCIATES**

**MON VALLEY-CHARLEROI**  
1200 McKean Avenue  
Charleroi, PA 15022  
724-483-1340 Phone

[www.WashingtonObGyn.net](http://www.WashingtonObGyn.net)

Because infection of the baby can cause serious problems, you may be treated during labor if you have any risk factors for beta strep infection, even if your test results are negative. The risk factors are:

- preterm labor (labor that starts before 37 weeks of pregnancy)
- premature rupture of membranes (water breaking before 37 weeks of pregnancy)
- membranes ruptured for longer than 18 hours
- a previous child with beta strep infection
- fever (higher than 100.4°F, or 38.0°C) during labor.
- GBS bacteria found sometime during the pregnancy.

Treatment with IV antibiotics during labor is usually very effective. Babies rarely develop the serious side effects of GBS infection with this treatment.

If a newborn develops any signs or symptoms of GBS infection, the baby is treated with IV antibiotics and watched very closely. The baby may stay in a special intensive care unit.

### **How long will the effects last?**

A woman who has GBS does not usually have any symptoms of illness or long-lasting effects. The main risk is that the newborn will become sick. GBS infection can affect the baby's blood, brain, spinal cord, or lungs. The baby may have long-term problems that affect hearing, vision, or learning abilities if the brain is infected. The baby might die.

### **How can I help prevent GBS infection of the baby?**

Getting tested for GBS and being treated with antibiotics during labor, if necessary, greatly reduces the risk that your baby will be infected with GBS.

Written by Edward Zabrek, MD.

Published by [RelayHealth](#).

Last modified: 2007-08-06

Last reviewed: 2007-07-30

This content is reviewed periodically and is subject to change as new health information becomes available. The information is intended to inform and educate and is not a replacement for medical evaluation, advice, diagnosis or treatment by a healthcare professional.

[Women's Health Advisor 2008.1 Index](#)

[Women's Health Advisor 2008.1 Credits](#)

© 2008 RelayHealth and/or one of its affiliates. All Rights Reserved.